

Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.

Prevalence of Post-traumatic Stress Disorder Among Survivors of Severe COVID-19 Infections: A Systematic Review and Meta-Analysis

Ramya Nagarajan, Yuvaraj Krishnamoorthy, Vanitha Basavarachar, Raghul Dakshinamoorthy


PII: S0165-0327(21)01262-3

DOI: https://doi.org/10.1016/j.jad.2021.11.040

Reference: JAD 14076

To appear in: Journal of Affective Disorders

Received date: 29 June 2021 Revised date: 13 October 2021 Accepted date: 14 November 2021



Please cite this article as: Ramya Nagarajan, Yuvaraj Krishnamoorthy, Vanitha Basavarachar, Raghul Dakshinamoorthy, Prevalence of Post-traumatic Stress Disorder Among Survivors of Severe COVID-19 Infections: A Systematic Review and Meta-Analysis, *Journal of Affective Disorders* (2021), doi: https://doi.org/10.1016/j.jad.2021.11.040

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2021 Published by Elsevier B.V.

**TITLE PAGE:** 

Prevalence of Post-traumatic Stress Disorder Among Survivors of Severe COVID-19

**Infections: A Systematic Review and Meta-Analysis** 

Authors:

Ramya Nagarajan, Yuvaraj Krishnamoorthy, Vanitha Basavarachar, Raghul

Dakshinamoorthy

Author affiliations:

Ramya Nagarajan, Scientist-B, National Institute of Epidemiology, Chennai, India

Yuvaraj Krishnamoorthy, Assistant Professor, Department of Community Medicine, ESIC

medical College and PGIMSR, Chennai, India

Vanitha Basavarachar, Senior Resident, Department of Community Medicine, Bowring and

Lady Curzon Medical College and Research Institute, Bengaluru, India

Raghul Dakshinamoorthy, Senior Resident, Department of Orthopedics, Institute of

medical Sciences, Banaras Hindu University, Varanasi, India

Corresponding Author.

Ramya Nagarajan, Consultant (India- EIS), National Institute of Epidemiology, Chennai,

India

Ph: +91 74834 02558

E.Mail: nagarajanramya55@gmail.com

Declaration of interests: None

1

### Highlights:

- 1. This is the first systematic review and meta-analysis to estimate the prevalence of PTSD among Severe cases of COVID-19 as per our knowledge.
- 2. The prevalence of PTSD among the severe cases of COVID-19 is high compared to the general public.
- 3. PTSD manifests as a post COVID symptom beyond four weeks of acquiring the COVID-19 infection. Hence resources must be allocated to follow up these patients closely.

#### **Abstract:**

**Background:** Post-traumatic stress disorder (PTSD) develops after a person has experienced a traumatic event which can be highly accounted for amidst the ongoing COVID-19 pandemic. This study aims to estimate the prevalence of PTSD among the severe cases of COVID-19.

**Methods:** We included the observational studies done to estimate the burden of PTSD among severe COVID-19 patients. Data was extracted manually using structured data extraction form and analyzed in STATA version 14.2. A random-effects model was applied, and the final pooled data was reported as proportion with a 95% confidence interval. Multivariable meta-regression analysis was carried out, and a forest plot was utilized to represent the study-specific and pooled estimates for overall and subgroup analysis.

**Results:** We included 13 articles with 1,093 participants in our analysis. The pooled prevalence was estimated to be 16% (95%CI: 9% to 23%). We found a substantial heterogeneity between the studies that reported the outcome ( $I^2=87.9\%$ , p<0.001). In subgroup analysis, the difference in prevalence estimate between the regions was statistically significant.

Limitations: We found significant between-study variability for the outcome. In addition,

our review was found to have substantial publication bias. We also found that the lower

quality of the majority of the studies being included in our review.

**Interpretation:** Our study states that the risk of PTSD is higher following severe COVID-19

infection. Understanding this burden will help us in diverting the resources and adapting

necessary interventions to control the situation.

Key Words: PTSD, COVID-19, Hospitalization, ICU admission

**Introduction:** 

A coronavirus is a group of viruses primarily affecting the respiratory tract. Most viruses that

cause human infections result in mild to moderate respiratory tract infections. COVID-19

was first notified to the WHO by the Chinese government in the second half of December

2019. Following the investigation, a novel coronavirus, SARS-CoV-2, was found to be the

causative factor. It later spread across the world.<sup>2</sup>

This pandemic has also faced various strict public health measures to limit transmission.

Some of these are regular use of outdoor masks, following physical distancing, adapting

handwashing techniques, and getting vaccinated against the disease.<sup>3,4</sup> Strict lockdown

measures were implemented in various countries globally, and most of the population was

advised to adapt to work from home. Many people have also lost their jobs in this process.

People who got infected were isolated from others while quarantining their contacts to avoid

disease transmission. This has also led to increased stigma in society. All these factors have

increased psychological morbidities in the general population.<sup>5</sup>

3

As per American Psychiatric Association, Posttraumatic stress disorder is a psychiatric disorder that may occur in people who have experienced or witnessed a traumatic event. Events such as a natural disaster, a severe accident, a terrorist act, war/combat, or rape or who have been threatened with death, sexual violence, or severe injury are factors associated with PTSD.<sup>6</sup> Many people exposed to a traumatic event experience symptoms of PTSD in the days following the event.<sup>6</sup>

Due to the disease, social stigma, prolonged hospital stays, economic loss, etc., some COVID-19 patients developed severe infections and required ICU admission during the treatment. These ICU admissions can be traumatic experiences for the patients. As a result, some might also develop the condition Post Traumatic Stress Disorder (PTSD). Hence, we conducted this systematic review and meta-analysis to estimate the prevalence of PTSD among the severe cases of COVID-19.

#### **Methods**

#### Design and Registration

We conducted this systematic review and meta-analysis by including the observational studies (cross-sectional, prospective, or retrospective studies) on the prevalence of PTSD among severe COVID-19 patients. The review was registered under the "International prospective register of systematic review (PROSPERO)" (PROSPERO ID: CRD42021256948). The latest 2020 checklist of Preferred Reporting Items for Systematic Review and Meta-Analysis (PRISMA) was used for reporting our review.<sup>7</sup>

### Eligibility criteria

#### Study design

We included observational studies (cross-sectional/prospective/retrospective studies) reporting the prevalence estimate of PTSD among severe COVID-19 patients. There wasn't

any restriction in age/study setting/country/region/any specific comorbidities. Studies were included without considering the mode of interview (online/offline). Abstracts, preprints, full-text articles met the inclusion criteria, while case reports, case series, and grey literature (unpublished data/articles) were excluded.

## Study participants

We used the following criteria to identify the studies conducted on severe COVID-19: patients with the respiratory rate (RR) > 30 breaths/min (or) oxygen saturation (SpO2) < 93% (or) oxygenation index (PaO2/FiO2)  $\le$  300 mmHg (or) requirement of ICU (or) mechanical ventilation. Studies must have followed up these patients after four weeks of COVID-19 diagnosis to be eligible for inclusion. Studies conducted among patients with only mild to moderate COVID-19 infection were excluded from the systematic review.

#### Outcome measures

We included the studies reporting the prevalence of PTSD irrespective of the scale used for measurement.

#### Search strategy

An extensive electronic database search was done in the following search engines:

MEDLINE, EMBASE, Google Scholar, and ScienceDirect. Search terms were built using

MESH terms in MEDLINE and EMTREE terms in EMBASE. The terms used to construct
the search strategy were 'Stress Disorders, Post-Traumatic' 'Posttraumatic stress Disorder'

Post Traumatic Stress Syndrome' 'COVID-19' 'hospitalization' 'severe COVID-19' 'ICU
admission.' Supplementary Appendix 1 shows the detailed search strategy used to extract
the studies for the review. The search was restricted from Jan 2020 to May 2021, and only
English-language articles were included.

#### Study selection

This process has involved three stages:

### Primary screening

Two independent investigators (RN & YK) have performed the preliminary screening of title, abstract, and keywords during the literature search. Full-text articles were retrieved for the studies relevant to the eligibility criteria.

#### Secondary screening

The two investigators (RN & YK) again screened the full text of these retrieved studies and assessed it against the review's eligibility criteria. Studies satisfying the eligibility criteria in terms of design, participants, and outcome were considered eligible to be included in our review.

## Finalizing the study selection

Disagreements during the primary and secondary screening process between the investigators were resolved by another investigator (VB). The final consensus on study inclusion was also reached with the help of the investigator (VB). "PRISMA flow chart" was utilized to represent the screening and selection process (**Figure 1**).

## Data extraction and management

Data was extracted manually by the primary investigator (RN). A structured data extraction form was developed and pilot-tested during the protocol stage itself. It consisted of the following information: general information (author, study title, publication year, country), details related to methods section (study design, study setting, study participants, sample size, diagnostic tool, and interview mode), and outcome (number of participants with PTSD).

The data was transferred by the primary investigator (RN) into the software STATA version 14.2. The data entered was double-checked by the second investigator (YK) to ensure correct entry by comparing the data presented in the review and the primary articles.

### Risk of Bias assessment

All the included studies were assessed by two authors (RN and YK) independently to look for the quality using the "Newcastle-Ottawa (NO) scale utilized for cross-sectional studies." It was assessed using two criteria (selection and outcome). Under selection criteria following domains were employed to evaluate the risk of bias: sample representativeness, sample size justification, rate of non-respondents and their characteristics, and use of validated measurement tool. To assess the risk of bias under outcome criteria, assessment of outcome through independent blind assessment or record linkage was used. Every outcome was rated as having high risk (1 point) or low risk (0 points) depending upon the quality of evidence and availability of information. An increased risk of bias was considered for studies with a score of  $\geq 3$  points.

### Statistical analysis

Meta-analysis was performed using the STATA 14.2 (StataCorp, College Station, TX, USA) software. Standard error was calculated using the reported number of outcomes and the sample size for each study. The "Metaprop" function in STATA was used for performing the pooled analysis. To minimize the effect of extremely small or large prevalence on the overall estimate, "Freeman Tukey double arcsine transformation" was done to stabilize the variance. Random-effects model was applied to account for heterogeneity between the included studies. The final pooled data was reported as proportion with a 95% confidence interval (CI).

Following methods were used to assess the evidence of heterogeneity: Chi-square test to identify the heterogeneity and I² statistic to quantify the heterogeneity. I² value < 25% considered to be mild, 25–75% moderate, and >75% was considered substantial heterogeneity. I² In our analysis, since all the included outcomes had significant heterogeneity, additional subgroup analysis, and meta-regression were performed. The potential covariates for performing meta-regression were geographical region, study design, measurement tool, mode of interview, mean age, sample size, and quality of the study. Multivariable meta-regression analysis included the study-level factors with p-value less than 0.20 in the univariable model.

Forest plot was used for graphical representation of the study-specific and pooled estimates for overall and subgroup analysis. Publication bias was checked and graphically represented by funnel plot and Doi plot. Asymmetry of the plot was tested using Egger's test and LFK index. P-value less than 0.10 in Egger's test is indicative of a possible publication bias. Based on the LFK index value, the possibility of publication bias is classified as no asymmetry (value within  $\pm 1$ ), minor asymmetry (value out of  $\pm 1$  but within  $\pm 2$ ), and significant asymmetry (value > $\pm 2$ ). <sup>13</sup>

#### Results

### Study Selection

During the primary screening, a total of 1597 records were obtained, in which 50 articles relevant to our review were extracted for full-text search. During the secondary screening, we went through the full-text of these articles and found 13 articles with 1,093 participants satisfying our inclusion criteria and included as a part of the analysis (**Figure 1**). <sup>14–26</sup>

### Characteristics of the studies included

Most of the studies were carried out in the European region (3 studies in Italy, two each in Netherlands and United Kingdom, each in France and Turkey). The remaining studies were carried out in China, Iran, and the United States of America. More than 50% of the studies (7 out of 13) conducted amidst the COVID-19 pandemic were cross-sectional, while the rest were prospective studies (cohort studies). The studies were undertaken after the patient got discharged from the hospital following admission for severe COVID-19. The follow-up or data collection time interval following discharge has ranged from 4 weeks to 16 weeks. The mean age of the participants ranges from 39.7 years to 60.9 years. The majority of the studies (8 out of 13) used PCL-5 (PTSD Checklist for DSM-5) to diagnose PTSD. In contrast, the rest of the studies used IES-R (Impact of Event Scale-Revised), CAPS-5 (Clinician-Administered PTSD scale for DSM-5), PTSS-14 (Post Traumatic Stress Syndrome), and TSQ (Trauma Screening Questionnaire) to diagnose PTSD among the severe COVID-19 patients. The interview mode was a telephonic interview (5 out of 13 studies), an offline interview by following up the patient in a hospital setting (5 out of 13 studies), and online applications in the rest of the studies. (Table 1).

### Risk of Bias assessment

All the studies had a high risk of bias concerning the representativeness of the sample and outcome assessment given the nature of data collection and mode of interview. In contrast, all the studies had a low risk of bias concerning using the validated tool. Only four studies reported a non-response rate and its characteristics, while the sample size justification is provided in most studies (8 out of 13). Finally, the review included 13 studies, out of which nine were of lower quality as per the NO scale (**Table 2**).

#### Prevalence of PTSD among severe COVID-19 patients

In total, 13 studies were included to find out the prevalence of PTSD among severe COVID-19 patients. The pooled prevalence was estimated to be 16% (95%CI: 9% to 23%) (**Figure 2**). We found substantial heterogeneity between the studies that reported the outcome (I<sup>2</sup>=87.9%, p<0.001). Subgroup analysis was carried out in terms of geographical region, study design, follow-up duration, measurement tool, and interview mode (**Table 3**).

#### Additional subgroup analysis

significant (Supplementary Figure 5).

Subgroup analysis based on the geographical region could not reveal much information. Most of the studies were carried out in the European region, and only one study was from the American, Western Pacific, and Eastern Mediterranean regions. However, the difference in prevalence estimate between the areas was statistically significant (p<0.001) (Supplementary Figure 1). Analysis based on study design revealed a higher pooled prevalence amongst the cross-sectional studies (19%) compared to cohort studies (13%). However, the difference was not statistically significant (Supplementary Figure 2). Based on data collection time point, the analysis showed that the prevalence of PTSD was similar irrespective of the time point or follow-up period (4 weeks period=18%; 12 weeks period=20%) (Supplementary Figure 3). Based on interview mode, the Subgroup analysis showed the highest pooled prevalence of PTSD (21%) for the studies conducted online. In comparison, the studies conducted through the offline interview (15%) and telephonic interview (14%) had almost similar estimates (Supplementary Figure 4). Analysis based on the assessment tool to diagnose PTSD revealed that the pooled prevalence of PTSD while using PCL-5 was 10% and 29% for the IES-R scale, and this difference was statistically

### Meta-regression

We also performed meta-regression to look for the heterogeneity sources with the study-level characteristics such as geographical region, study design, measurement tool, mode of interview, mean age, sample size, and quality of the study. Neither of the factors was significantly explaining the effect estimate. However, multivariable meta-regression analyses were performed with factors with a p-value less than 0.20, such as sample size, geographical region, measurement tool, and interview mode. Though the adjusted model could explain 100% of the heterogeneity, it was not statistically significant (**Table 4**).

### Additional analysis

Publication bias was graphically assessed by funnel plot (**Supplementary Figure 6**) and Doi plot (**Supplementary Figure 7**). Both the plots showed signs of asymmetry, statistically confirmed by Egger's test (p=0.07) and the LFK index (3.18). Sensitivity analysis was carried out to check the robustness of the estimates. It has shown no significant variation in the magnitude or direction of the outcome, indicating that a single study failed to influence the overall pooled estimate (**Supplementary Figure 8**).

#### **Discussion**

PTSD can occur in people who experience a traumatic event in their lifetime.<sup>6</sup> The risk of increased prevalence of PTSD has also been observed in previous coronavirus pandemics, making its occurrence during this COVID-19 pandemic highly explainable.<sup>27</sup> Some of the severe cases of COVID-19 result in mortality. The fear of death might be among the many reasons responsible for PTSD amongst these patients. The death of any close relative in the family may also lead to the occurrence of PTSD. Therefore, it is necessary to understand the prevalence of PTSD amongst these patients to highlight its importance in every aspect of health during such testing times.

Our review showed that 16% of the severe COVID-19 patients globally had PTSD.

Geographic distribution of the prevalence did not reveal any additional information due to limited studies in regions other than Europe. The evidence is limited on the post mental effects of severely COVID-19 affected individuals. The estimated prevalence of PTSD in our study was higher than the existing literature focussing on all the COVID-19 patients irrespective of the severity. However, the prevalence was comparatively lower compared to previous similar pandemics. <sup>28</sup>

A meta-analysis on the survivors of ICU-admitted patients with SARS and MERS infection has revealed that about 39% of them had suffered from PTSD. There could be plenty of reasons for such contrasting differences in the prevalence across these studies. However, some critical reasons were; high lethality and lower epidemic spread of the SARS and MERS compared to COVID-19, the difference in the follow-up period for PTSD, measurement tool, study region, study quality across these reviews. However, a large-scale web-based survey conducted by the Center for Disease Control and Prevention (CDC) among 5000 US adults showed that almost all suffered from at least one adverse mental or behavioral health problem concerned with this pandemic. The survey also reported that more than one-fourth of them had symptoms of trauma and stress-related disorder. This was substantially higher than the lifetime prevalence of PTSD (6.8%) suffered by the US adults as stated by the National Institute of Mental Health of Americas. Variations in the geographical distribution of PTSD explain the severity of COVID-19 hit zones. Such variations in mental disorders should occur as they take a more significant toll on individuals exposing them to vulnerability as well as the methodological differences in the individual studies.

These findings further necessitate the need for having a more significant number of longitudinal studies assessing the prevalence of PTSD amongst severe patients in heavily affected countries in Asia and America like India, the USA, Brazil, etc. PTSD being a public

health problem, should be identified and treated at the earliest. It is essential to follow up with these patients after getting discharged from the hospital for severe COVID-19 infection. The patient often reports somatic symptoms, but the psychological symptoms go unnoticed unless a healthcare worker follows up. Therefore, it is recommended to consider PTSD when the patients come for post-COVID follow-up.

The major strength of our review was the rigorous literature search and methodology followed to provide reliable estimates. Additionally, this was also the first review reporting the prevalence of PTSD amongst severe COVID-19 patients. We have also carried out additional subgroup analysis, meta-regression, and sensitivity analysis to find the source of heterogeneity and check the robustness of the results. However, our review has certain limitations. We found significant between-study variability (significant chi-square test for heterogeneity and I² statistics) for the outcome. For such high heterogeneity, the methodological differences between the included studies can be held accountable, such as type of patients categorized as severe COVID-19, geographical region, sample size, tools used for diagnosis, quality of the studies, etc. In addition, our review was found to have substantial publication bias. We also found that the lower quality of the majority of the studies included in our review might further limit the generalisability of our study findings.

Irrespective of the limitations seen in our review, the findings did show some important implications for clinical management. PTSD is a psychiatric disorder in individuals who have experienced severe life-threatening incidences, war/ combat, rape, serious trauma. It affects all age groups and both genders. In our case, the COVID-19 pandemic has been threatening the lives of millions of people, which makes it a traumatic event by itself irrespective of suffering from the disease. On top of this, hospitalization with severe COVID-19 infection or admission in ICU under critical condition might further amplify the risk associated with PTSD. Apart from the disease, hospitalized patients also face other difficulties such as

financial prevalence pertaining to the treatment, staying away from home, stigma in society,

etc. This meta-analysis helps identify this vital aspect and urges clinicians and psychiatrists

worldwide to provide extra attention and care for patients admitted or discharged following

severe COVID-19 conditions.

Though our results provide some crucial information for better understanding the prevalence

of PTSD following severe COVID-19, it is still needed to perform a substantial number of

high-quality longitudinal studies to establish the prevalence of PTSD more precisely.

Nevertheless, understanding this prevalence will help us divert the resources and adapt

necessary interventions to control the situation.

**Acknowledgment:** Nil

**Contributions:** Nagarajan R, acquisition of data, analysis, and interpretation of data, drafting

the article, final approval; Krishnamoorthy Y, conception and design of the study, acquisition

of data, analysis, and interpretation of data, drafting the article, final approval; Basavarachar

V revising the article, final approval: Dakshinamoorthy R, critical revision, final approval.

Funding: This research did not receive any specific grant from funding agencies in the

public, commercial, or not-for-profit sectors.

**Declaration of interests**: None

**References:** 

1. Zheng J. SARS-CoV-2: an Emerging Coronavirus that Causes a Global Threat.

2020;16.

2. World Health Organization. Timeline: WHO's COVID-19 response [Internet]. World

Health Organization. 2021. Available from:

https://www.who.int/emergencies/diseases/novel-coronavirus-2019/interactive-

14

timeline

- 3. Ministry of Health and Family Welfare [Internet]. Government of India. 2021.

  Available from: https://www.mohfw.gov.in/
- World Health Organization. Coronavirus Disease (COVID-19) Pandemic [Internet].
   2021. Available from: https://www.who.int/emergencies/diseases/novel-coronavirus-
- 5. Krishnamoorthy Y, Nagarajan R, Kumar G, Menon V. Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information.
  2020;(January).
- 6. Felix Torres. What Is Posttraumatic Stress Disorder? [Internet]. Americal Psychiatric Association. 2020. Available from: https://www.psychiatry.org/patients-families/ptsd/what-is-ptsd
- 7. Title TT. PRISMA 2020 Checklist PRISMA 2020 Checklist. 2020;2020–1.
- 8. Cascella M, Rajnik M, Aleem A, Dulebohn SC, Di Napoli R. Features, Evaluation, and Treatment of Coronavirus (COVID-19) [Internet]. StatPearls. 2021. Available from: http://www.ncbi.nlm.nih.gov/pubmed/32150360
- 9. Post-COVID Conditions [Internet]. Centre for Disease Control. 2021. Available from: https://www.cdc.gov/coronavirus/2019-ncov/long-term-effects.html
- 10. Shea B, Robertson J, Peterson J, Welch V, Losos M. The Newcastle-Ottawa Scale ( NOS ) for Assessing the Quality of Nonrandomized Studies in Meta- Analysis Bias and Confounding Newcastle-Ottowa Scale.

- 11. Nyaga VN, Arbyn M, Aerts M. M E T H O D O LO G Y O pen A ccess Metaprop : a Stata command to perform meta-analysis of binomial data. 2014;1–10.
- 12. Higgins JPT, Thomas J, Chandler J, Cumpston M, Li T, Page MJ WV. Cochrane
  Handbook for Systematic Reviews of Interventions [Internet]. Higgins JPT, Thomas J,
  Chandler J, Cumpston M, Li T, Page MJ WV, editor. Cochrane; 2021. Available from:
  www.training.cochrane.org/handbook.
- 13. Egger M, Smith GD, Schneider M, Minder C. Bias in meta-analysis detected by a simple, graphical test. 1997;315(September).
- Janiri D, Carfì A, Kotzalidis GD, Bernabei R, Landi F, Sani G. Posttraumatic Stress
   Disorder in Patients After Severe COVID-19 Infection. JAMA psychiatry. 2021
   May;78(5):567–9.
- Morin L, Savale L, Pham T, Colle R, Figueiredo S, Harrois A, et al. Four-Month
   Clinical Status of a Cohort of Patients after Hospitalization for COVID-19. JAMA J
   Am Med Assoc. 2021;325(15):1525–34.
- 16. Naidu SB, Shah AJ, Saigal A, Smith C, Brill SE, Goldring J, et al. The high mental health burden of "Long COVID" and its association with on-going physical and respiratory symptoms in all adults discharged from hospital. The European respiratory journal. 2021.
- 17. Rass V, Beer R, Schiefecker AJ, Kofler M, Lindner A, Mahlknecht P, et al.

  Neurological outcome and quality of life 3 months after COVID-19: A prospective observational cohort study. Eur J Neurol. 2021 Mar;
- 18. Khademi M, Vaziri-Harami R, Shams J. Prevalence of Mental Health Problems and Its Associated Factors Among Recovered COVID-19 Patients During the Pandemic: A

- Single-Center Study. Front psychiatry. 2021;12:602244.
- 19. Liu D, Baumeister RF, Veilleux JC, Chen C, Liu W, Yue Y, et al. Risk factors associated with mental illness in hospital discharged patients infected with COVID-19 in Wuhan, China. Psychiatry Res [Internet]. 2020 Oct;292(January):113297. Available from: https://linkinghub.elsevier.com/retrieve/pii/S0165178120317601
- 20. de Graaf MA, Antoni ML, ter Kuile MM, Arbous MS, Duinisveld AJF, Feltkamp MCW, et al. Short-term outpatient follow-up of COVID-19 patients: A multidisciplinary approach. EClinicalMedicine. 2021;32.
- 21. Mazza MG, De Lorenzo R, Conte C, Poletti S, Vai B, Bollettini I, et al. Anxiety and depression in COVID-19 survivors: Role of inflammatory and clinical predictors.
  Brain Behav Immun [Internet]. 2020 Oct;89(January):594–600. Available from: doi: 10.1016/j.bbi.2020.07.037
- 22. Martillo MA, Dangayach NS, Tabacof L, Spielman LA, Dams-O'Connor K, Chan CC, et al. Postintensive Care Syndrome in Survivors of Critical Illness Related to Coronavirus Disease 2019. Crit Care Med. 2021;Publish Ah:1–12.
- 23. Parker AJ, Humbir A, Tiwary P, Mishra M, Shanmugam M, Bhatia K, et al. Recovery after critical illness in COVID-19 ICU survivors. Br J Anaesth [Internet]. 2021;126(6):e217–9. Available from: https://doi.org/10.1016/j.bja.2021.03.005
- 24. Beck K, Vincent A, Becker C, Keller A, Cam H, Schaefert R, et al. Prevalence and factors associated with psychological burden in COVID-19 patients and their relatives: A prospective observational cohort study. PLoS One [Internet]. 2021;16(5 May):1–30. Available from: http://dx.doi.org/10.1371/journal.pone.0250590
- 25. Tarsitani L, Vassalini P, Koukopoulos A, Borrazzo C, Alessi F, Di Nicolantonio C, et

- al. Post-traumatic Stress Disorder Among COVID-19 Survivors at 3-Month Follow-up After Hospital Discharge. J Gen Intern Med. 2021 Mar;1–6.
- 26. Poyraz BÇ, Poyraz CA, Olgun Y, Gürel Ö, Alkan S, Özdemir YE, et al. Psychiatric morbidity and protracted symptoms after COVID-19. Psychiatry Res. 2021 Jan;295:113604.
- 27. Yuan K, Gong YM, Liu L, Sun YK, Tian SS, Wang YJ, et al. Prevalence of posttraumatic stress disorder after infectious disease pandemics in the twenty-first century, including COVID-19: a meta-analysis and systematic review. Mol Psychiatry [Internet]. 2021; Available from: http://dx.doi.org/10.1038/s41380-021-01036-x
- 28. Ahmed H, Patel K, Greenwood DC, Halpin S, Lewthwaite P, Salawu A, et al. Long-term clinical outcomes in survivors of severe acute respiratory syndrome (SARS) and Middle East respiratory syndrome (MERS) coronavirus outbreaks after hospitalisation or ICU admission: A systematic review and meta-analysis. J Rehabil Med. 2020;52(5).
- 29. Dietze PM, Peacock A, Mariottini C, Ojanperä I, Kriikku P, Wang QQ, et al. Mental Health, Substance Use, and Suicidal Ideation During the COVID-19. Addict Behav [Internet]. 2020;29(32):1049–57. Available from: https://doi.org/10.1016/j.addbeh.2020.106754%0Ahttp://dx.doi.org/10.1038/s41380-020-00830-7
- 30. Post Traumatic Stress Disorder [Internet]. National Institute of Mental Health. 2017.

  Available from: https://www.nimh.nih.gov/health/statistics/post-traumatic-stress-disorder-ptsd
- 31. Torres F. What Is Posttraumatic Stress Disorder? Americal Psychiatric Association.2020.

Table 1: Characteristics of the included studies (N=13)

S.	Author and	Country	Study design	Sample Size	PTSD	Follow up	Assessme	Severity Criteria	Mode of	Mean Age
No.	Year				among	duration	nt tool		interview	(in years)
					SCP					
1.	Beck 2021	Netherlands	Cohort study	Overall: 126	2	4 weeks	IES-R	ICU admissions	Telephone	58.2
				SCP: 18		1				
2.	Graff 2020	Netherlands	Cross-sectional	Overall: 81	2	6 weeks	PCL-5	ICU admissions	Offline	60.8
			study	SCP: 34						
3.	Janiri 2021	Italy	Cross-sectional	Overall: 381	23	4 weeks	CAPS-5	ICU admissions	Offline	55.26
3.	Jann 1 2021	Italy	study	SCP: 65	23	4 WCCKS	CAI 5-5	ico admissions	Offinic	33.20
			study	SCI . 05						
4.	Khademi	Iran	Cross-sectional	Overall: 602	16	4 weeks	PCL-5	Hospitalization	Telephone	53.2
	2021		study	SCP: 418				with severe		
								illness		
5.	Liu 2020	China	Cross-sectional	Overall: 675	5	4 weeks	PCL-5	ICU admissions	Online	53.58
			study	SCP: 35						
6.	Martillo	USA	Cohort study	Overall: 45	8	4 weeks	PCL-5	ICU admissions	Telephone	53.9
	2021									

7.	Mazza 2020	Italy	Cohort study	Overall:402 SCP: 220	40	4 weeks	PCL-5	Hospitalization with Severe Illness	Offline	57.8
8.	Morin 2021	France	Cohort study	Overall: 117 SCP: 94	7	16 weeks	PCL-5	ICU admissions	Offline	60.9
9.	Naidu 2021	United Kingdom	Cross-sectional study	Overall: 760 SCP: 90	12	8 weeks	TSQ	ICU admissions	Online	57
10.	Parker 2021	United Kingdom	Cross-sectional study	Overall: 36 SCP: 17	8	12 weeks	PTSS-14	ICU admissions	Telephone	52.5
11.	Poyraz 2021	Turkey	Cross-sectional study	Overall: 284 SCP: 34	14	4 weeks	IES-R	Hospitalization with Severe Illness	Online	39.7
12.	Rass 2021	Austria	Cohort study	Overall: 135 SCP: 31	4	12 weeks	PCL-5	ICU admissions	Offline	56
13.	Tarsitani 2021	Italy	Cohort study	Overall: 115 SCP: 26	2	12 weeks	PCL-5	ICU admissions	Telephone	57

ICU: Intensive Care Unit

PTSD: Post Traumatic Stress Disorder

IES-R: Impact of Event Scale-Revised

PCL-5: PTSD Checklist for DMS-5

CAPS-5: Clinician-Administered PTSD Scale for DMS-5

TSQ: Trauma Screening Questionnaire- PTSD

PTSS-14: Post Traumatic Stress syndrome-14

USA: United States of America

SCP: Severe COVID-19 Patients

Table 2: Risk of Bias Assessment with Newcastle Ottawa Scale (N=13)

S.No	Study	Representativeness of	Justification of	Non-response rate	Use of a validated	Outcome	Overall
		the sample	sample size		tool	assessment	Quality
1.	Beck 2021	High	Low	Low	Low	High	High
2.	Graff 2020	High	Low	High	Low	High	Low
3.	Janiri 2021	High	High	High	Low	High	Low
4.	Khademi 2021	High	Low	Low	Low	High	High
5.	Liu 2020	High	High	High	Low	High	Low
6.	Martillo 2021	High	High	High	Low	High	Low
7.	Mazza 2020	High	Low	High	Low	High	Low
8.	Morin 2021	High	Low	Low	Low	High	High
9.	Naidu 2021	High	High	High	Low	High	Low
10.	Parker 2021	High	Low	High	Low	High	Low
11.	Poyraz 2021	Hìgh	Low	High	Low	High	Low
12.	Rass 2021	High	Low	Low	Low	High	High
13.	Tarsitani 2021	High	High	High	Low	High	Low

Table 3: Summary of findings and subgroup analysis of studies reporting the prevalence of PTSD among severe COVID-19 patients (N=13)

Characteristics	Number of studies pooled	Pooled proportion (95% CI)
Pooled Prevalence of PTS	S among sever COVID-19 patie	nts= 16% (95%CI: 9%- 23%)
Geographical regions		
Europe	10	18% (11%-26%)
America	1	18% (8%-32%)
Western Pacific	1	14% (5%-30%)
Eastern Mediterranean	1	4% (2%- 6%)
Study design	<sup>4</sup> C	
Cross sectional study	7	19% (7%- 37%)
Cohort study	6	13% (8%-18%)
Follow-up duration		
4 weeks	7	18% (8%-31%)
6 weeks	1	6% (1%- 20%)
8 weeks	1	13% (7%- 22%)
12 weeks	3	20% (3%-43%)
16 weeks	1	7% (3%- 15%)
The tool used to diagnose	PTSD	
PCL-5	8	10% (5%- 17%)
IES-R	2	29% (17%-42%)
PTSS-14	1	47% (23% - 72%)
TSQ	1	13% (7%-22%)
CAPS-5	1	35% (24%- 48%)

### **Mode of interview**

Offline	5	15% (7%-25%)
Telephonic Interview	5	14% (3%- 30%)
Online	3	21% (8%- 37%)

IES-R: Impact of Event Scale- Revised

PCL-5: PTSD Checklist for DMS-5

CAPS-5: Clinician Administered PTSD Scale for DMS-5

TSQ: Trauma Screening Questionnaire- PTSD

PTSS-14: Post Traumatic Stress syndrome-14

Table 4: Univariable and multivariable meta-regression results (N=13)

Characteristics	Unadjusted	Unadjusted p-	Adjusted	Adjusted p-				
	Coefficient	value	Coefficient	Value				
Sample Size	-0.0003	0.052	0.0005	0.462				
Geographical region								
Europe	Ref	-	Ref	-				
Americas	-0.003	0.985	0.08	0.75				
Western Pacific	-0.03	0.831	-0.20	0.64				
Eastern	-0.14	0.050	-0.27	0.49				
Mediterranean		4	0					
Diagnostic tool		0)	•					
PCL-5	Ref		Ref	-				
IES-R	0.19	0.20	0.03	0.90				
PTSS-14	0.37	0.16	0.39	0.32				
TSQ	0.03	0.76	-0.24	0.56				
CAPS-5	0.25	0.09	0.26	0.20				
Mode of Interview								
Offline	Ref	-	Ref	-				
Telephonic	-0.10	0.14	0.01	0.95				
Online	0.02	0.81	0.28	0.51				

### Figure captions:

Figure 1: PRISMA Flowchart




Figure 2: Forest plot showing the pooled prevalence of PTSD among severe COVID-19 patients (N=13)

